

Reche-Soto, P.; Cardona, D.; Díaz, A.; Gómez-Carmona, C.; Pino-Ortega, J. (201x) ACELT y PLAYERLOAD™: Dos variables para la cuantificación de la carga neuromuscular / ACELT and PLAYERLOAD™: Two Variables to Quantify Neuromuscular Load. Revista Internacional de Medicina y Ciencias de la Actividad Física y el Deporte vol. X (X) pp. xx. [Http://cdeporte.rediris.es/revista/___*](http://cdeporte.rediris.es/revista/)

ORIGINAL

ACELT AND PLAYERLOAD™: TWO VARIABLES TO QUANTIFY NEUROMUSCULAR LOAD

ACELT Y PLAYERLOAD™: DOS VARIABLES PARA LA CUANTIFICACIÓN DE LA CARGA NEUROMUSCULAR

Reche-Soto, P.¹; Cardona, D.²; Díaz, A.³; Gómez-Carmona, C.⁴ y Pino-Ortega, J.⁵

¹ PhD student in Sports Science. Faculty of Sports Science and Physical Activity, University of Murcia (Spain) pedrorechesto@gmail.com

² Doctor of Sports Science. Professor at the Faculty of Physical Education, Recreation and Sports, Colombian Polytechnic Jaime Isaza Cadavid (Colombia) donaldpf@gmail.com

³ Research Group Ingesport. Professor and Full Professor in the Faculty of Sports Science and Physical Activity, University of Murcia (Spain) ardiaz@um.es

⁴ Training Optimization and Sports Performance Group (GOERD). PhD in Sports Science. Faculty of Sports Science and Physical Activity, University of Extremadura (Spain) cdgomezcarmona@unex.es

⁵ Department of Sport and Physical Activity. Doctor and Full Professor in the Faculty of Sports Science and Physical Activity, University of Murcia (Spain) josepinoortega@um.es

Spanish-English revision: Diana Schofield Smith, schofielddiane@gmail.com

FINANCING

The author Carlos D. Gómez Carmona is a beneficiary of a grant from the Ministry of Education, Culture and Sport (FPU17 / 00407). This investigation has not received funding from any public, private or non-profit agent.

CONFLICT OF INTERESTS

The last author of this article is an advisor in the area of Sports Science in the company responsible for the development of the inertial device used. To guarantee the objectivity of the results, this author has not contributed to the data analysis nor to the results section, but he has contributed significantly to other parts of the manuscript. The authors certify that this study has not been published or is in the process of being considered for publication in another journal, accepting the publication rules of the Journal.

Código UNESCO / UNESCO code: 5899 Otras especialidades pedagógicas (Educación Física y Deporte). / Other pedagogical specialties (Physical Education and Sport)

Clasificación Consejo de Europa / Council of Europe Classification: Otras (Rendimiento deportivo) / Other (sport performance).

Recibido 17 de marzo de 2018 **Received** March 17, 2018

Aceptado 15 de enero de 2019 **Accepted** January 15, 2019

ABSTRACT

The objectives of this study are: (i) to describe the dynamics of the PlayerLoad™ and AcelT variables and (ii) to quantify the neuromuscular load at different anatomical points (back, centre of mass, knee and ankle) during an incremental treadmill ramp test. Twenty-three semi-professional national football players participated voluntarily in this study (age: 22.56 ± 4.8 years, body mass: 75.5 ± 5.5 kg, height: 1.79 ± 0.5 m). Both variables were recorded using four WIMU PRO™ inertial devices. The main results indicate that: (1) there is a nearly perfect correlation between both variables ($r = 0.931$), (2) the highest values in both variables have been found in the knee ($PL^{\text{TM}} = 8.01 \pm 2.76$; $AccelT = 2.70 \pm 0.50$) and the ankle ($PL^{\text{TM}} = 7.85 \pm 2.27$, $AccelT = 2.87 \pm 0.49$) and (3) there is wide inter-subject variability. In conclusion, PlayerLoad™ and AcelT are two valid indicators for the analysis and quantification of neuromuscular demands.

KEYWORDS: accelerometry, football, performance, AcelT, PlayerLoad™.

RESUMEN

Los objetivos de esta investigación son: (i) describir el comportamiento de las variables PlayerLoad™ y AcelT y (ii) cuantificar la carga neuromuscular en diferentes puntos anatómicos (espalda, centro de masas, rodilla y tobillo) durante un test incremental en rampa en tapiz rodante. Veintitrés jugadores semiprofesionales de fútbol varones de categoría nacional participaron voluntariamente en este estudio (edad: $22,56 \pm 4,8$ años; masa corporal: $75,5 \pm 5,5$ kg; altura: $1,79 \pm 0,5$ m). Ambas variables se registraron empleando 4 dispositivos inerciales WIMU PRO™. Los principales resultados encontrados indican que: (1) existe una correlación casi perfecta entre ambas variables ($r=0,931$), (2) los mayores valores en ambas variables se han encontrado en la rodilla ($PL^{\text{TM}} = 8,01 \pm 2,76$; $AccelT = 2,70 \pm 0,50$) y el tobillo ($PL^{\text{TM}} = 7,85 \pm 2,27$; $AccelT = 2,87 \pm 0,49$) y (3) existe una amplia variabilidad intersujeto. En conclusión, PlayerLoad™ y AcelT son dos indicadores válidos para el análisis y cuantificación de las demandas neuromusculares.

PALABRAS CLAVE: acelerometría, fútbol, rendimiento, AcelT, PlayerLoad™.

1. INTRODUCTION

Among outdoor team sports, football is one of the most popular, both in the number of players and fans. This aspect has caused this sport modality to be one of the most studied in scientific literature (Sánchez, Yagüe, Fernández and Petisco, 2014). The current problem is that the great complexity of this sport has meant that the research carried out has not completely reduced the

uncertainty surrounding training strategies for improving performance (Aguiar, Botelho, Lago, Maças and Sampaio, 2012).

In terms of physical and physiological demands, football is considered a team sport involving intermittent efforts of high intensity (McMillan, Helgerud, Macdonal and Hoff, 2005), in which a large number of short sprints, rapid accelerations, decelerations, turns, jumps, kicks and tackles are performed. Thus, a full recovery between actions in many cases is not possible (Arnason et al, 2004). All these dynamic and unpredictable technical-tactical actions, which vary in duration and intensity (Bloomfield, Polman and O'Donoghue, 2007), constitute the total internal and external load that players experience (Akubat, Barrett and Abt, 2014) and pose an energy challenge for them (Stølen, Chamari, Castagna and Wisløff, 2005). Accurate and objective quantification of the activities performed by players is fundamental to understanding the physical demands of football (Bradley, Di Mascio, Peart and Olsen, 2010, Dwyer and Gabbett, 2012, Johnston et al., 2012), since up to 1,350 different actions performed by the players have been recorded (Mohr, Krstrup and Bangsbo, 2003).

In order to improve the preparation methods, it is necessary to expand scientific knowledge. Therefore, an objective analysis of the characteristics of the game in competitive situations is required (Carling, Williams and Reilly, 2006). This possibility is, in large part, enabled by technological development and the authorisation of FIFA (International Federation of Association Football), for the use of electronic devices in competitive situations (FIFA, 2018). These investigations are providing new evidence in the study of physical and tactical demands (Castellano and Casamichana, 2014). Being aware of the requirements of the competition allows us to control training loads more effectively (Grehaigne, Godbout, and Zerai, 2011) and to improve the design of training tasks both physically (Gómez-Carmona, Gamonales, Pino-Ortega and Ibáñez, 2018) and tactically (Reche-Soto, Cardona, Díaz, Gómez-Carmona and Pino-Ortega, 2018b).

Monitoring the loads of team sports players is a common practice in both training and competition (Rogalski, Dawson, Heasman and Gabbett, 2013). The recording of the parameters is used by sports professionals in order to provide an explanation for possible changes in the performance of the athlete or to try to reduce the risk of injury, illness or training overload (Halson, 2014). The heterogeneous nature of each athlete's response to this training load makes their individualised analysis fundamental (Brink, Nederhof, Visscher, Schmikli and Lemmink, 2010, Paulson, Mason, Rhodes and Goosey-Tolfrey, 2015).

Considering the nature of the workload, it is classified as internal or external, both in the sports and scientific fields (Bartlett, O'Connor, Pitchford, Torres-Ronda and Robertson, 2017, Costa et al., 2013). Internal load refers to the psychological and physiological stress resulting from training-competition, while the external load includes the work done by the athlete in terms of distance, speed and acceleration (Lambert and Borresen, 2010; McLaren, Weston, Smith, Cramb and Portas, 2016; Paulson et al., 2015). In team sports, there is greater difficulty in evaluating the internal load (Borresen and Lambert, 2009),

which has meant that the quantification of external load has undergone greater development in recent years.

Accelerometers were introduced in the 1950s (Culhane, O'Connor, Lyons, and Lyons 2005), and have evolved technologically to offer sufficient quality and reliability with a low production cost. They allow us to quantitatively evaluate human movement with a fully portable device. (Lemoine, Coroian, Mastroianni and Grundfest, 2008). The multifunctional conception of inertial devices (Inertial measurement units, IMUs) is increasingly used in the sports context. These devices include different sensors such as accelerometers, gyroscopes, magnetometers, GNSS, etc. (Akenhead and Nassis, 2016; Boyd, Ball and Aughey, 2013; Gabbett, 2015).

One of the most commonly used load indicators, based on the accelerometer signal, is the PlayerLoad™ (PL) variable (Barrett et al., 2016, Bradley et al., 2010, Dalen, Jørgen, Gertjan, Havard and Ulrik, 2016; Reche-Soto et al., 2018a), also known as Body Load (Gomez-Piriz, Jiménez-Reyes and Ruiz-Ruiz, 2011), depending on the inertial device used. This indicator is the combination of the accelerations produced in the three anatomical planes of movement, producing an estimate of the total load (Cummins, Orr, O'Connor and West, 2013), expressed in arbitrary units (Barrett et al., 2016). The PL™ has been proved to be a reliable and valid indicator (Hollville, Couturier, Guilhem and Rabita, 2016), which has a high correlation with physiological variables such as heart rate and VO2max (Barrett et al., 2016), and subjective scales of perception of effort (Casamichana, Castellano, Calleja-Gonzalez, San Román and Castagna, 2013). In addition, this variable has obtained high inter- and intra-device test-retest reliability in cyclical (Barrett et al., 2016) and acyclic activities (Boyd et al., 2013) as well as in multidirectional tasks (ICC = 0.806-0.949) (Barreira et al., 2017).

For all of the above, the objectives of this study were: (i) to describe the dynamics of the neuromuscular load variables PlayerLoad™ and AceIT in an incremental treadmill ramp test and (ii) to quantify the neuromuscular load at different anatomical points (back, lumbar area, knee and ankle) during this test.

2. METHOD

2.1. PARTICIPANTS

Twenty-three semi-professional football players from a national football league (Third Division, Group XIII) voluntarily participated in this study (Age: 22.56 ± 4.8 years, Weight: 75.5 ± 5.5 kg, Height: 1.79 ± 0.5 metres). All participants had to meet the following requirements: (i) have more than two years of experience in football practice at a national level, (ii) present more than one year of experience with advanced technological monitoring in both training and official matches and (iii) not present any musculoskeletal injury or health problems that prevented their evaluation.

The study was approved by the ethics committee of the University of Murcia before the start (registration number 2061/2018), in accordance with the code of ethics of the World Medical Association (Declaration of Helsinki, 2013). Participants were previously informed of the details of the investigation and its possible risks and benefits, and gave their informed consent.

2.2. VARIABLES

• **PlayerLoad™**: it is the vector sum of the accelerations of the device in the 3 axes (vertical, anteroposterior and lateral). This variable has been used to evaluate the neuromuscular load in different athletes (Cormack, Mooney, Morgan and McGuigan, 2013) and is calculated from the following equation:

$$Player\ Load = \frac{\sqrt{(a_{y1} - a_{y-1})^2 + (a_{x1} - a_{x-1})^2 + (a_{z1} - a_{z-1})^2}}{100}$$

• **AccelT**: it is the magnitude of acceleration or resulting vector (Waldron, Twist, Highton, Worsfold, and Daniels, 2011) identified as the vector sum of the total acceleration recorded by the accelerometer, which is the result of gravity (y axis), changes in horizontal movement (x axis) and forces related to the rotation movements (z-axis) of a body segment or object to which the accelerometer is attached (O'Donovan, Kamnik, O'Keeffe and Lyons, 2007, Kunze, Bahle, Lukowicz and Partridge, 2010). It is calculated from the following formula:

$$Resultant\ vector\ (AccelT) = \sqrt{x^2 + y^2 + z^2}$$

2.3. INSTRUMENTS

Anthropometric characteristics

The height of the subjects was measured with a wall-mounted height meter during full inhalation (SECA, Hamburg, Germany), and body weight was obtained using a body composition monitor, model BC-601 (TANITA, Tokyo, Japan).

Neuromuscular load

Both variables were recorded using 4 WIMUPRO™ inertial devices (RealTrack Systems, Almeria, Spain), which contain four triaxle accelerometers that detect and measure movement using an electromechanical system with a sampling frequency from 10 to 1,000 Hz. The motion detection range is $\pm 16\text{ g}$, $\pm 16\text{ g}$, $\pm 32\text{ g}$ and $\pm 400\text{ g}$. Each device has its own 1 GHz microprocessor, 8 GB of internal memory and high-speed USB output to record, store and download data. Each device has an internal battery with a duration of more than 4 hours, weighing 70 g and measuring 81 × 45 × 16 mm. In this study, the accelerometers were configured to record the research variables with a sampling frequency of 1,000 Hz.

The devices were placed on the: (i) back (C6, between the scapulae), (ii) lumbar region (L3, at the height of the centre of mass) (McGregor, Armstrong, Yaggie, Parshad and Bollt, 2011), (iii) knee (5 cm above the crack of the kneecap) and (iv) ankle (5 cm above the lateral malleolus). The devices were placed on the outside part of the right leg in all subjects, both on the ankle and knee.

Prior to their placement, the devices were calibrated and synchronised. Thanks to this process, the accelerometers eliminate the 4 sources of error that may occur: displacement error, scaling error, orthogonal errors and random error (Wang, Liu and Fan, 2006). The device calibration process was carried out following the manufacturer's recommendations in the auto-start process. To ensure proper functioning, three aspects had to be met: (a) leaving the device immobile for 30 seconds, (b) on a flat surface and (c) without close contact with magnetic devices (Bastida-Castillo, Gómez-Carmona, Reche, Granero-Gil and Pino-Ortega, 2018). Following this procedure, the accelerometers in this device have obtained very high reliability values in static and dynamic tests in different anatomical locations (Gómez-Carmona, Bastida-Castillo, García-Rubio, Ibáñez and Pino-Ortega, 2018).

Figure 1 shows the location of the inertial devices, which were placed as follows: (i) back, using a specific harness anatomically adjusted to the subject and (ii) L3, ankle and knee, using an extendable band that was attached to the subject with Velcro and finally sealed with adhesive tape in order to reinforce the fastening.



Figure 1. Location of the inertial devices in the study participants.

2.4. PROCEDURE

The study was carried over 2 weeks, with a total of 3 sessions. In the first session, anthropometric measurements were recorded and both the protocol and the objective of the study were explained. The second session consisted of familiarisation with the test and running using technological monitoring. In the third session, the incremental ramp test was performed on a treadmill. The starting speed was 8 km / h. From the beginning, a continuous speed increase of 0.1 km / h was made every 12 seconds (1 km / h every 2 minutes). The test ended when the subject could not maintain the speed.

All sessions began at 9:00 am and the subjects had to meet the following requirements: (i) suppression of alcohol and caffeine intake 24 hours before each session and (ii) not performing high intensity physical activity the 48 hours prior to the completion of the protocol; so that none of these factors interfered in the investigation (Billat, 2002, Spriet, 2014).

Before carrying out each of the protocols, a standardised warm-up of 5 minutes of continuous running was performed at 65% of HRmax, which was monitored and controlled in real time by sending data from the WIMUPRO™ inertial devices (RealTrack Systems, Almeria, Spain) using Wi-Fi technology to a computer which had the software SPRO™ (RealTrack Systems, Almeria, Spain) to check the perfect functioning of the devices (Bastida-Castillo, Gómez-Carmona and Pino-Ortega, 2016). At the end of the protocol, subjects performed 5 minutes of continuous running at 55% of the HRmax.

3. STATISTICAL ANALYSIS

The Shapiro-Wilk test was used to determine the distribution of the data, and for the homoscedasticity the Levene test was applied (Field, 2013). The analysis showed a normal distribution, so parametric tests were used. First a descriptive analysis was made showing the results in terms of average and standard deviation. A linear regression was performed to determine the cause-effect relationship between the two neuromuscular load variables PlayerLoad™ and AcelT depending on the location of the inertial devices, the subjects involved and the running speed. The statistical tests were performed using the software SPSS 24.0 (SPSS Inc., Chicago IL, USA). The statistical significance was established with the value of $p < 0.05$.

4. RESULTS

Table 1 presents the descriptive analysis of the neuromuscular load variables AcelT and PlayerLoad™ in the different locations and according to the different running speeds. Higher values are found in both variables the higher the running speed. In addition, values also increase as the location is closer to the point of contact of the foot with the ground.

Table 1. Descriptive analysis of neuromuscular load variables.

Speed (km/h)	PlayerLoad™								AcelT							
	Back		L3		Knee		Ankle		Back		L3		Knee		Ankle	
	M	SD	M	SD	M	SD	M	SD	M	SD	M	SD	M	SD	M	SD
8	2.26	0.35	2.37	0.42	4.90	0.63	4.81	0.82	1.19	0.04	1.30	0.06	2.05	0.09	2.13	0.10
8,5	2.34	0.40	2.51	0.43	5.26	0.67	5.12	0.67	1.20	0.04	1.32	0.06	2.14	0.09	2.22	0.10
9	2.46	0.43	2.66	0.42	5.65	0.70	5.60	0.70	1.22	0.04	1.34	0.06	2.22	0.09	2.33	0.09
9,5	2.51	0.44	2.80	0.44	6.02	0.77	5.97	0.68	1.23	0.04	1.36	0.07	2.30	0.10	2.43	0.10
10	2.60	0.44	2.94	0.46	6.45	0.82	6.38	0.73	1.25	0.04	1.38	0.07	2.39	0.10	2.53	0.09
10,5	2.70	0.46	3.08	0.47	6.80	0.90	6.74	0.75	1.26	0.04	1.40	0.08	2.46	0.11	2.62	0.10
11	2.80	0.45	3.22	0.46	7.44	1.55	7.08	0.83	1.27	0.04	1.42	0.08	2.59	0.22	2.72	0.10
11,5	2.91	0.44	3.34	0.45	7.91	1.79	7.57	0.83	1.28	0.04	1.44	0.08	2.68	0.25	2.82	0.10
12	3.00	0.47	3.49	0.44	8.32	1.93	7.96	0.83	1.29	0.04	1.45	0.08	2.76	0.27	2.92	0.10
12,5	3.11	0.46	3.63	0.47	8.76	2.08	8.46	0.84	1.30	0.04	1.47	0.09	2.85	0.30	3.04	0.11
13	3.21	0.48	3.72	0.46	9.25	2.13	8.93	0.93	1.32	0.04	1.48	0.09	2.94	0.31	3.15	0.12
13,5	3.30	0.46	3.87	0.47	9.78	2.29	9.46	1.04	1.33	0.04	1.50	0.08	3.05	0.36	3.25	0.13
14	3.47	0.51	3.93	0.43	10.53	3.00	9.99	1.06	1.34	0.04	1.51	0.08	3.22	0.53	3.36	0.12
14,5	3.55	0.49	4.09	0.44	11.40	4.00	10.54	1.16	1.36	0.05	1.52	0.08	3.37	0.67	3.47	0.12
15	3.56	0.47	4.16	0.46	10.82	1.79	11.23	1.26	1.37	0.05	1.54	0.09	3.24	0.12	3.61	0.12
15,5	3.75	0.44	4.40	0.52	10.93	1.41	11.79	1.21	1.39	0.06	1.57	0.09	3.31	0.11	3.72	0.14
16	3.73	0.30	4.58	0.27	11.68	0.78	12.68	1.06	1.37	0.03	1.59	0.12	3.44	0.04	3.94	0.07
Total	2.94	0.63	3.37	0.75	8.01	2.76	7.85	2.27	1.28	0.07	1.43	0.11	2.70	0.50	2.87	0.49

Nota. M: mean; SD: Standard deviation.

Figure 2 shows the regression analysis between the variables PlayerLoad™ and AcelT during the incremental running test in all the subjects analysed. There are different dynamics between the upper body load and the lower body load along the speed spectrum. As a whole, there is a nearly perfect correlation between both neuromuscular load variables ($R^2 = 0.931$).




Figure 2. Graphical representation of the relationship between the variables PlayerLoad™ and AcelT during the progressive incremental test on the treadmill.

Table 2 shows the linear regression between the neuromuscular load variables AcelT and PlayerLoad™ in each of the subjects analysed in the different locations. We find an inter-subject variability in the location analysis. In addition, we find the best relationships in the ankle ($R^2 = 0.956$) and the knee ($R^2 = 0.916$) both individually and globally. Both locations belong to the neuromuscular load dynamics of the lower body.

Table 2. Linear regression between the variables of neuromuscular load PlayerLoad™ and AcelT in the different locations analysed in each of the study participants.

Participants	Locations			
	Ankle	Knee	L3	Back
1	0.997	0.958	0.998	0.973
2	0.982	0.996	0.980	0.970
3	0.992	0.993	0.992	0.947
4	0.992	0.993	0.992	0.947
5	0.990	0.998	0.980	0.941
6	0.997	0.999	0.990	0.964
7	0.996	0.984	0.992	0.967
8	0.993	0.990	0.973	0.989
9	0.996	0.996	0.988	0.956
10	0.996	0.998	0.984	0.923
11	0.987	0.996	0.954	0.988
12	0.998	0.993	0.987	0.976
13	0.993	0.979	0.997	0.987
14	0.998	0.999	0.997	0.988
15	0.990	0.998	0.997	0.986
16	0.992	0.997	0.956	0.955
17	0.995	0.988	0.956	0.952
18	0.987	0.987	0.949	0.990
19	0.990	0.994	0.962	0.978
20	0.989	0.998	0.956	0.987
21	0.996	0.992	0.993	0.952
22	0.995	0.998	0.998	0.887
23	0.995	0.996	0.980	0.973
Total	0.956	0.916	0.765	0.713

Finally, Table 3 presents a relational analysis of the neuromuscular load variables PlayerLoad™ and AcelT in the different locations according to the running speed. Better relationships are found when the speeds are higher in the aforementioned locations, ankle and knee.

Table 3. Linear regression between the neuromuscular load variables PlayerLoadTM and AceIT in the different locations analysed according to the running speed.

Speed	Locations				Total
	Ankle	Knee	L3	Back	
8	0.806	0.628	0.520	0.139	0.879
8.5	0.606	0.646	0.514	0.276	0.900
9	0.590	0.577	0.531	0.331	0.910
9.5	0.545	0.478	0.555	0.361	0.913
10	0.535	0.440	0.551	0.424	0.916
10.5	0.549	0.407	0.559	0.429	0.916
11	0.765	0.789	0.513	0.469	0.906
11.5	0.789	0.834	0.524	0.421	0.909
12	0.847	0.834	0.543	0.394	0.908
12.5	0.820	0.847	0.563	0.464	0.912
13	0.817	0.841	0.566	0.550	0.915
13.5	0.817	0.835	0.462	0.462	0.921
14	0.907	0.904	0.420	0.546	0.935
14.5	0.928	0.920	0.396	0.495	0.924
15	0.786	0.676	0.379	0.494	0.928
15.5	0.780	0.608	0.541	0.512	0.933
16	0.781	0.897	0.567	0.549	0.972

5. DISCUSSION

The objectives of this study were: (i) to describe the dynamics of the neuromuscular load variables PlayerLoadTM and AceIT and (ii) to quantify such demands in different anatomical locations along the speed spectrum during an incremental treadmill ramp test.

In the descriptive analysis performed, the highest values in the PlayerLoadTM and AceIT variables were found in the knee (PLTM = 8.01 ± 2.76, AceIT = 2.70 ± 0.50) and the ankle (PLTM = 7, 85 ± 2.27, AceIT = 2.87 ± 0.49), the values in both variables increasing as the running speed increased. The contribution of the load from individual planes can also be influenced by the anatomical position of the accelerometer. Barrett, Midgley and Lovell (2014) find that the values of both variables in the locations which are closest to the point of contact of the foot with the ground increase as the speed increases. However, it is generally accepted that the centre of mass (lumbar area, L3) (McGregor et al., 2011) is the optimal anatomical location for the placement of these devices (Halsey, Shepard and Wilson, 2011; McGregor, Busa, Yaggie and Bollt, 2009), although, as in our case, there are exceptions in the literature (Boyd et al., 2013; Scott, Black, Quinn and Coutts, 2013; Cormack et al., 2013). Specifically, in the study conducted by Barrett et al., (2014) where the registered PLTM was slightly higher in L3 than in the back.

As for the regression analysis performed between the accelerometric load variables PlayerLoadTM and AceIT, it is shown that both variables have a nearly perfect correlation ($r > 0.931$). Therefore, both variables are valid for the quantification and analysis of neuromuscular demands. The PLTM has been accepted as a valid indicator to interpret the amount of external load experienced by a player in different sports such as football (Barreira et al., 2017; Casamichana et al., 2013; Scott et al., 2013), Australian football (Boyd et

al., 2013; Gastin, McLean, Spittle and Breed, 2013; Scott et al., 2013) and linear running (Barrett et al., 2014). Previous research has shown significant and strong relationships between neuromuscular load measurements and the ratio of injuries in rugby players (Gabbett, 2004a, Gabbett 2004b), so both variables are put forward as indicators of injury risk in team sports. The PL™ also presents a high relation with respect to the Edwards index ($r = 0.70$), the subjective perception of effort (PSE) ($r = 0.74$) and with the volume of movement through the total distance travelled variable ($r = 0.70$) (Casamichana et al., 2013). Scott et al. (2013) also found a strong relationship between the PL™ variable and the Edwards index ($r = 0.80$). In short, the studies show that the PL™ is a valid indicator to quantify the demands in football, being acceptable for use in competition ($CV = 1.9\%$) (Boyd et al., 2013).

The linear regression between the neuromuscular load variables AcelT and PlayerLoad™ in each of the subjects analysed in the different locations shows an inter-subject variability in the location analysis. In addition, we found the best relationships in the ankle ($R^2 = 0,956$) and knee ($R^2 = 0,916$), both individually and globally. Such relationships are higher at a higher running speed. The results obtained are linked to those found in the research conducted by Nedergaard et al. (2017) where there is a low correlation between the acceleration of the centre of mass (L3) and the rest of locations (tibia, knee and scapula) concluding that the acceleration of the whole body cannot be detected by placing an accelerometer in the centre of mass due to the complexity of sports actions. Therefore, to achieve the best accuracy, the ideal location would be the ankle / tibia because it is the first joint point that receives more directly the forces that the subject exerts against the ground while running, obtaining a high validity with respect to force platforms (Raper et al., 2018) and detecting differences in impact depending on the type of footwear (Sinclair and Sant, 2017).

6. LIMITATIONS OF THE STUDY

Different limitations must be taken into account when interpreting the results obtained in this investigation. Firstly, the number of participants is small ($n = 23$), which may influence the statistical power of the results analysed. Secondly, the participants analysed were national level male football players, so the results could not be extrapolated to other study populations. Finally, only four inertial devices at a specific sampling frequency were used for the data collection. The components of the inertial devices, the calibration of the sensors and the sampling frequency can affect the results obtained. Therefore, the recording of the data through the inertial devices was carried out following the manufacturer's recommendations.

7. CONCLUSIONS

PlayerLoad™ and the AcelT are two valid indicators for the analysis and quantification of neuromuscular demands. Therefore, both indicators can be used interchangeably to quantify neuromuscular load in training and competition. The lower limb supports greater neuromuscular load with respect to the upper limb, therefore, it is necessary to analyse its dynamics to adapt

training loads and recovery protocols. Finally, there is wide inter-subject variability, which is why an individualised load analysis is recommended for greater specificity with the aim of improving sports performance.

8. REFERENCES

- Aguiar, M.; Botelho, G.; Lago, C.; Maças, V., & Sampaio, J. (2012). A review on the effects of soccer small-sided games. *Journal of human kinetics*, 33, 103–113.
- Akenhead, R., & Nassis, G. P. (2016). Training load and player monitoring in high-level football: current practice and perceptions. *International journal of sports physiology and performance*, 11(5), 587-593.
- Akubat, I., Barrett, S., & Abt, G. (2014). Integrating the internal and external training loads in soccer. *International journal of sports physiology and performance*, 9(3), 457-462.
- Arnason, A., Sigurdsson, S. B., Gudmundsson, A., Holme, I., Engebretsen, L., & Bahr, R. (2004). Physical fitness, injuries, and team performance in soccer. *Medicine & Science in Sports & Exercise*, 36(2), 278-285.
- Barreira, P., Robinson, M. A., Drust, B., Nedergaard, N., Raja Azidin, R. M. F., & Vanrenterghem, J. (2017). Mechanical Player Load™ using trunk-mounted accelerometry in football: Is it a reliable, task-and player-specific observation? *Journal of Sports Sciences*, 35(17), 1674-1681
- Barrett, S., Midgley, A. W., Towlson, C., Garrett, A., Portas, M., & Lovell, R. (2016). Within-match playerload™ patterns during a simulated soccer match: Potential implications for unit positioning and fatigue management. *International Journal of Sports Physiology and Performance*, 11(1), 135-140
- Barrett, S., Midgley, A., & Lovell, R. (2014). PlayerLoad™: reliability, convergent validity, and influence of unit position during treadmill running. *International journal of sports physiology and performance*, 9(6), 945-952.
- Bartlett, J. D., O'Connor, F., Pitchford, N., Torres-Ronda, L., & Robertson, S. J. (2017). Relationships between internal and external training load in team-sport athletes: evidence for an individualized approach. *International Journal of Sports Physiology and Performance*, 12(2), 230-234
- Bastida-Castillo, A., Gómez-Carmona, C. D., y Pino-Ortega, J. (2016). Efectos del Tipo de Recuperación Sobre la Oxigenación Muscular Durante el Ejercicio de Sentadilla. *Kronos*, 15(2).
- Bastida-Castillo, A., Gómez-Carmona, C. D., Reche, P., Granero-Gil, P. y Pino-Ortega, J. (2018). Valoración de la estabilidad del tronco mediante un dispositivo inercial. *Retos: nuevas tendencias en educación física, deporte y recreación*, (33), 199-203.
- Billat, V. (2002). *Fisiología y metodología del entrenamiento. De la teoría a la práctica* (Vol. 24). Editorial Paidotribo: España.
- Bloomfield, J., Polman, R., & O'Donoghue, P. (2007). Physical demands of different positions in FA Premier League soccer. *J Sports Sci Med*, 6(1), 63-70.
- Borresen, J., & Lambert, M. I. (2009). The quantification of training load, the training response and the effect on performance. *Sports Medicine*, 39(9),

779-795.

- Boyd, L. J., Ball, K., & Aughey, R. J. (2013). Quantifying external load in Australian football matches and training using accelerometers. *International journal of sports physiology and performance*, 8(1), 44-51.
- Bradley, P. S., Di Mascio, M., Peart, D., Olsen, P., & Sheldon, B. (2010). High-intensity activity profiles of elite soccer players at different performance levels. *The Journal of Strength & Conditioning Research*, 24(9), 2343-2351.
- Brink, M. S., Nederhof, E., Visscher, C., Schmikli, S. L., & Lemmink, K. A. (2010). Monitoring load, recovery, and performance in young elite soccer players. *The Journal of Strength & Conditioning Research*, 24(3), 597-603.
- Carling, C., Williams, A. M., & Reilly, T. (2006). Handbook of soccer match analysis: a systematic approach to improving performance. *Journal of sports science & medicine*, 5(1), 171.
- Casamichana, D., Castellano, J., Calleja-Gonzalez, J., San Román, J., & Castagna, C. (2013). Relationship between indicators of training load in soccer players. *The Journal of Strength & Conditioning Research*, 27(2), 369-374.
- Castellano, J., y Casamichana, D. (2014). Deporte con dispositivos de posicionamiento global (GPS): aplicaciones y limitaciones. *Revista de Psicología del Deporte*, 23(2), 355-364.
- Cormack, S. J., Mooney, M. G., Morgan, W., & McGuigan, M. R. (2013). Influence of neuromuscular fatigue on accelerometer load in elite Australian football players. *International journal of sports physiology and performance*, 8(4), 373-378.
- Costa, E. C., Vieira, C. M., Moreira, A., Ugrinowitsch, C., Castagna, C., & Aoki, M. S. (2013). Monitoring external and internal loads of Brazilian soccer referees during official matches. *Journal of Sports Science & Medicine*, 12(3), 559-564.
- Culhane, K. M., O'connor, M., Lyons, D., & Lyons, G. M. (2005). Accelerometers in rehabilitation medicine for older adults. *Age and ageing*, 34(6), 556-560.
- Cummins, C., Orr, R., O'Connor, H., & West, C. (2013). Global positioning systems (GPS) and microtechnology sensors in team sports: a systematic review. *Sports Medicine*, 43(10), 1025-1042.
- Dalen, T., Jørgen, I., Gertjan, E., Havard, H. G., & Ulrik, W. (2016). Player Load, Acceleration, and Deceleration During Forty-Five Competitive Matches of Elite Soccer. *The Journal of Strength & Conditioning Research*, 30(2), 351-359.
- Dwyer, D. B., & Gabbett, T. J. (2012). Global positioning system data analysis: velocity ranges and a new definition of sprinting for field sport athletes. *The Journal of Strength & Conditioning Research*, 26(3), 818-824.
- Field, A. (2013). *Discovering statistics using IBM SPSS statistics*. Sage: London.
- FIFA (2017). Regla 4. El equipamiento de los jugadores. In: FIFA, ed., *Reglas de Juego 2017-2018*, 1st ed. Zurich, p.57.

- Gabbett, T. J. (2004a). Influence of training and match intensity on injuries in rugby league. *Journal of Sports Sciences*, 22, 409– 417.
- Gabbett, T. J. (2004b). Reductions in pre-season training loads reduce injury rates in rugby league players. *British journal of sports medicine*, 38(6), 743-749.
- Gabbett, T. J. (2015). Relationship between accelerometer load, collisions, and repeated high-intensity effort activity in rugby league players. *The Journal of Strength & Conditioning Research*, 29(12), 3424-3431.
- Gastin, P. B., McLean, O., Spittle, M., & Breed, R. V. (2013). Quantification of tackling demands in professional Australian football using integrated wearable athlete tracking technology. *Journal of science and medicine in sport*, 16(6), 589-593.
- Gómez-Carmona, C. D., Bastida-Castillo, A., García-Rubio, J., Ibáñez, S. J., & Pino-Ortega, J. (2018). Static and dynamic reliability of WIMU PRO™ accelerometers according to anatomical placement. *Proceedings of the Institution of Mechanical Engineers, Part P: Journal of Sports Engineering and Technology*, 1754337118816922.
- Gómez-Carmona, C., Gamonales, J., Pino-Ortega, J., & Ibáñez, S. (2018). Comparative Analysis of Load Profile between Small-Sided Games and Official Matches in Youth Soccer Players. *Sports*, 6, 173.
- Gomez-Piriz, P. T., Jiménez-Reyes, P., & Ruiz-Ruiz, C. (2011). Relation between total body load and session rating of perceived exertion in professional soccer players. *The Journal of Strength & Conditioning Research*, 25(8), 2100-2103
- Grehaigne, J. F., Godbout, P., & Zerai, Z. (2011). How the “rapport de forces” evolves in a soccer match: the dynamics of collective decisions in a complex system. *Revista de Psicología del Deporte*, 20(2), 747-765.
- Halsey, L. G., Shepard, E. L., & Wilson, R. P. (2011). Assessing the development and application of the accelerometry technique for estimating energy expenditure. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 158(3), 305-314.
- Halson, S. L. (2014). Monitoring training load to understand fatigue in athletes. *Sports Medicine*, 44(2), 139-147
- Hollville, E., Couturier, A., Guilhem, G., & Rabita, G. (2016, May). MinimaxX player load as an index of the center of mass displacement? A validation study. In *ISBS-Conference Proceedings Archive* (Vol. 33, No. 1)
- Johnston, R. J., Watsford, M. L., Pine, M. J., Spurr, R. W., Murphy, A. J., & Pruyn, E. C. (2012). The validity and reliability of 5-Hz global positioning system units to measure team sport movement demands. *The Journal of Strength & Conditioning Research*, 26(3), 758-765.
- Kunze, K., Bahle, G., Lukowicz, P., & Partridge, K. (2010, October). Can magnetic field sensors replace gyroscopes in wearable sensing applications? In *Wearable Computers (ISWC), 2010 International Symposium on* (pp. 1-4). IEEE.
- Lambert, M. I., & Borresen, J. (2010). Measuring training load in sports. *International Journal of Sports physiology and performance*, 5(3), 406-411
- Lemoyne, R., Coroian, C., Mastroianni, T., & Grundfest, W. (2008). Accelerometers for quantification of gait and movement disorders: a

- perspective review. *Journal of Mechanics in Medicine and Biology*, 8(02), 137-152.
- Mallo, J., & Navarro, E. (2008). Physical load imposed on soccer players during small-sided training games. *Journal of sports medicine and physical fitness*, 48(2), 166-171.
- McGregor, S. J., Armstrong, W. J., Yaggie, J. A., Bollt, E. M., Parshad, R., Bailey, J. J., ... & Kelly, S. R. (2011). Lower extremity fatigue increases complexity of postural control during a single-legged stance. *Journal of neuroengineering and rehabilitation*, 8(1), 43.
- McGregor, S. J., Busa, M. A., Yaggie, J. A., & Bollt, E. M. (2009). High resolution MEMS accelerometers to estimate VO₂ and compare running mechanics between highly trained inter-collegiate and untrained runners. *PLoS One*, 4(10), e7355.
- McLaren, S. J., Weston, M., Smith, A., Cramb, R., & Portas, M. D. (2016). Variability of physical performance and player match loads in professional rugby union. *Journal of science and medicine in sport*, 19(6), 493-497.
- McMillan, K., Helgerud, J., Macdonald, R., & Hoff, J. (2005). Physiological adaptations to soccer specific endurance training in professional youth soccer players. *British journal of sports medicine*, 39(5), 273-277.
- Mohr, M., Krstrup, P., & Bangsbo, J. (2003). Match performance of high-standard soccer players with special reference to development of fatigue. *Journal of sports sciences*, 21(7), 519-528.
- Nedergaard, N. J., Robinson, M. A., Eusterwiemann, E., Drust, B., Lisboa, P. J., & Vanrenterghem, J. (2017). The Relationship Between Whole-Body External Loading and Body-Worn Accelerometry During Team-Sport Movements. *International journal of sports physiology and performance*, 12(1), 18-26.
- O'Donovan, K. J., Kamnik, R., O'Keeffe, D. T., & Lyons, G. M. (2007). An inertial and magnetic sensor based technique for joint angle measurement. *Journal of biomechanics*, 40(12), 2604-2611.
- Paulson, T. A., Mason, B., Rhodes, J., & Goosey-Tolfrey, V. L. (2015). Individualized internal and external training load relationships in elite wheelchair rugby players. *Frontiers in physiology*, 6, e388.
- Raper, D. P., Witchalls, J., Philips, E. J., Knight, E., Drew, M. K., & Waddington, G. (2018). Use of a tibial accelerometer to measure ground reaction force in running: A reliability and validity comparison with force plates. *Journal of science and medicine in sport*, 21(1), 84-88.
- Reche-Soto, P., Cardona, D., Díaz, A., Gómez-Carmona, C. D., & Pino-Ortega, J. (2018b). Tactical Demands of Small-Sided Games in Football: Influence of Tracking Technology. *Revista Internacional de Medicina y Ciencias de la Actividad Física y el Deporte*, in press.
- Reche-Soto, P., Cardona-Nieto, D., Diaz-Suarez, A., Bastida-Castillo, A., Gomez-Carmona, C., Garcia-Rubio, J., & Pino-Ortega, J. (2018a). Player Load and Metabolic Power Dynamics as Load Quantifiers in Soccer. *Journal of Human Kinetics*, in press.
- Rogalski, B., Dawson, B., Heasman, J., & Gabbett, T. J. (2013). Training and game loads and injury risk in elite Australian footballers. *Journal of Science and Medicine in Sport*, 16(6), 499-503.

- Sánchez-Sánchez, J., Yagüe, J. M., Fernández, R. C., & Petisco, C. (2014). Efectos de un entrenamiento con juegos reducidos sobre la técnica y la condición física de jóvenes futbolistas. *RICYDE. Revista Internacional de Ciencias del Deporte*, 10(37), 221-234.
- Scott, T. J., Black, C. R., Quinn, J., & Coutts, A. J. (2013). Validity and reliability of the session-RPE method for quantifying training in Australian football: a comparison of the CR10 and CR100 scales. *The Journal of Strength & Conditioning Research*, 27(1), 270-276.
- Sinclair, J., & Sant, B. (2017). The effects of cross-fit footwear on the kinetics and kinematics of running. *Footwear Science*, 9(1), 41-48.
- Sports Medicine, 38, 743–749.
- Spriet, L. L. (2014). Exercise and sport performance with low doses of caffeine. *Sports medicine*, 44(2), 175-184
- Stølen, T., Chamari, K., Castagna, C., & Wisløff, U. (2005). Physiology of soccer. *Sports medicine*, 35(6), 501-536.
- Waldron, M., Twist, C., Highton, J., Worsfold, P., & Daniels, M. (2011). Movement and physiological match demands of elite rugby league using portable global positioning systems. *Journal of Sports Sciences*, 29(11), 1223-1230.
- Wang, J., Liu, Y., & Fan, W. (2006, June). Design and calibration for a smart inertial measurement unit for autonomous helicopters using MEMS sensors. *In Mechatronics and Automation, Proceedings of the 2006 IEEE International Conference on* (pp. 956-961). IEEE.

Número de citas totales / Total references: 63 (100%)

Número de citas propias de la revista / Journal's own references: 2 (3,17%)